

Polarized structure functions of the deuteron

S. Kumano *

Department of Physics
Saga University
Saga 840-8502, Japan

Talk given at the 14th International
Spin Physics Symposium

Osaka, Japan, October 16 – 21, 2000
(talk on Oct. 16, 2000)

* Email: kumanos@cc.saga-u.ac.jp. Information on his research is available at
<http://www-hs.phys.saga-u.ac.jp>.

Polarized structure functions of the deuteron

S. Kumano

Department of Physics, Saga University, Saga 840-8502, Japan

Abstract. Physics of spin-1 hadron is an unexplored topic in the high-energy region although spin-1/2 physics has been well investigated in the last decade. It is important to test our knowledge of hadron spin structure in a quite different field of spin physics. We discuss tensor structure functions, which do not exist for the spin-1/2 nucleon, in lepton scattering and in hadron reactions such as the polarized proton-deuteron Drell-Yan process.

INTRODUCTION

High-energy spin physics is entering into a new era in the sense that the difference between a naive quark-model prediction and experimental data is now roughly understood and that much detailed studies are in progress with the completion of the RHIC facility. So far, spin-structure studies have been focused on the spin-1/2 nucleon. For higher-spin hadrons, we know that different spin physics exists such as the tensor structure in the deuteron. In order to test our understanding of the hadron structure, it is important to investigate other spin quantities. In this sense, spin-1 hadrons, for example the deuteron, should be suitable for future investigations.

In this paper, the present status is explained for the deuteron spin structure in the high-energy region. First, polarized electron-deuteron scattering is discussed briefly in connection with the deuteron spin physics. In particular, new polarized structure functions are introduced. Then, as an alternative method, the polarized proton-deuteron Drell-Yan process is discussed for finding the new distributions.

SPIN-1 STRUCTURE IN ELECTRON SCATTERING

A general formalism of the deep inelastic electron-deuteron scattering is discussed in Ref. [1]. It suggests that there exist additional structure functions in comparison with the electron-proton scattering due to the spin-1 nature of the deuteron. We find that there are eight independent amplitudes for $\gamma(h_1) + d(H_1) \rightarrow \gamma(h_2) + d(H_2)$, where $h_{1,2}$ and $H_{1,2}$ are helicities, by using momentum conservation, parity

invariance, and time-reversal invariance. Four of them are usual structure functions: F_1 , F_2 , g_1 , and g_2 , which exist in the spin-1/2 proton. In addition, there are four new structure functions: b_1 , b_2 , b_3 , and b_4 . The b_3 and b_4 are higher-twist functions and b_2 is related to b_1 by the “Callan-Gross type” relation $b_2 = 2xb_1$ in the Bjorken scaling limit, so that the most essential part of new physics is contained in b_1 . A phenomenological sum rule $\int dx b_1(x) = 0$ was proposed [2] if there is no tensor polarization for antiquark distributions; however, it does not mean that $b_1(x)$ itself vanishes. This sum rule is similar to the Gottfried sum rule in a way [3]. Because there is no solid reason why the antiquark tensor polarization vanishes, the violation of the above sum rule could suggest a finite tensor polarization as the Gottfried-sum-rule violation indicated a finite \bar{u}/\bar{d} asymmetry.

There are three twist-2 structure functions: F_1 , g_1 , and b_1 , and they are related to electron-deuteron scattering cross sections as

$$\begin{aligned} F_1 &\propto \langle d\sigma \rangle , \\ g_1 &\propto d\sigma(\uparrow, +1) - d\sigma(\uparrow, -1) , \\ b_1 &\propto d\sigma(0) - [d\sigma(+1) + d\sigma(-1)]/2 . \end{aligned} \tag{1}$$

Here, $d\sigma(\uparrow, +1)$ and $d\sigma(\uparrow, -1)$ indicate spin parallel and anti-parallel cross sections, respectively. On the other hand, $\sigma(0)$ and $d\sigma(\pm 1)$ indicate cross sections with the deuteron spin state 0 and ± 1 . This fact means that the electron does not have to be polarized for measuring b_1 . There are a number of predictions on the x dependence of b_1 , but we cannot discuss the model results because of the space limitation. The interested reader may find a list of papers in the reference section of Ref. [4]. At this stage, b_1 is expected to be quite small, typically $b_1/F_1 \sim$ a few percent or much less. However, our experience of g_1 implies that the predictions could be wrong. In particular, the tensor structure in the high-energy region is still an unexamined topic experimentally. The HERMES experiment for measuring b_1 is in progress and its results will be reported in the near future. Therefore, the b_1 physics could become one of popular topics in hadron physics.

PROTON-DEUTERON DRELL-YAN PROCESS

In addition to the lepton scattering studies, it is, in principle, possible to investigate the same spin physics in hadron facilities. However, there are only a few theoretical studies on the spin-1 physics in hadron collisions. So far, the polarized proton-deuteron (pd) Drell-Yan reaction has been investigated in connection with the tensor structure of the deuteron [4,5].

Imposing the conditions of Hermiticity, parity conservation, and time-reversal invariance, we found many structure functions in the pd Drell-Yan. Even after integrating the cross section over the transverse momentum of the virtual photon \vec{Q}_T or after taking the limit $Q_T \rightarrow 0$, there are 22 functions [4]:

$$\begin{aligned}
\text{unpolarized :} & \quad W_{0,0}, W_{2,0} \\
\text{L/T polarized :} & \quad V_{0,0}^{LL}, V_{0,0}^{TT}, V_{2,0}^{LL}, V_{2,0}^{TT}, U_{2,1}^{TU}, U_{2,1}^{UT}, U_{2,1}^{TL}, U_{2,1}^{LT}, U_{2,2}^{TT} \\
\text{tensor polarized :} & \quad V_{0,0}^{UQ_0}, V_{0,0}^{TQ_1}, V_{2,0}^{UQ_0}, V_{2,0}^{TQ_1}, U_{2,1}^{TQ_0}, U_{2,1}^{LQ_1}, U_{2,1}^{TQ_2}, U_{2,1}^{UQ_1}, \\
& \quad U_{2,2}^{UQ_2}, U_{2,2}^{TQ_1}, U_{2,2}^{LQ_2}.
\end{aligned} \tag{2}$$

The subscripts of $F_{L,M}^{pol_p pol_d}$ ($F=W, V$, or U) indicate that it is obtained by the integration $\int d\Omega Y_{LM} d\sigma/(d^4Q d\Omega)$. The superscripts indicate the polarization states of the proton and deuteron. The notations U , L , and T indicate unpolarized, longitudinally polarized, and transversely polarized states. The quadrupole polarizations Q_0 , Q_1 , and Q_2 are specific in the reactions with a spin-1 hadron, and they are associated with the quadrupole terms: Q_0 for the term $3 \cos^2 \beta_B - 1 \sim Y_{20}$, Q_1 for $\sin \beta_B \cos \beta_B \sim Y_{21}$, and Q_2 for $\sin^2 \beta_B \sim Y_{22}$ with the polarization angle of the deuteron β_B . The structure functions in the categories “unpolarized” and “L/T polarized” exist in the proton-proton (pp) Drell-Yan. The new functions are listed as “tensor polarized” and they are associated with the spin-1 deuteron.

In defining the spin asymmetry, we should be careful about the tensor contributions. If it is defined in the usual way, for example $A_{LL}^{usual} = [\sigma(\uparrow, -1) - \sigma(\uparrow, +1)]/[\sigma(\uparrow, -1) + \sigma(\uparrow, +1)]$, the denominator is not equal to the unpolarized cross section because of

$$2 \langle \sigma \rangle = \sigma(\uparrow, +1) + \sigma(\uparrow, -1) + \frac{1}{3} [2 \sigma(\uparrow, 0) - \sigma(\uparrow, +1) - \sigma(\uparrow, -1)]. \tag{3}$$

In order to avoid the above tensor contribution, the unpolarized cross section is used for the denominator in Ref. [4] as a “theoretical” definition of the spin asymmetry. Using the obtained pd cross section with the structure functions in Eq.(2), we find the following spin asymmetries:

$$\begin{aligned}
& \langle \sigma \rangle, A_{LL}, A_{TT}, A_{LT}, A_{TL}, A_{UT}, A_{TU}, \\
& A_{UQ_0}, A_{TQ_0}, A_{UQ_1}, A_{LQ_1}, A_{TQ_1}, A_{UQ_2}, A_{LQ_2}, A_{TQ_2}.
\end{aligned} \tag{4}$$

Those in the first line exist in the pp reaction. However, the ones in the second line are specific for the pd Drell-Yan, and they are related to the tensor polarizations Q_0 , Q_1 , and Q_2 . For example, A_{UQ_0} , A_{UQ_1} , and A_{UQ_2} are given as

$$\begin{aligned}
A_{UQ_0} &= \frac{1}{2 \langle \sigma \rangle} \left[\sigma(\bullet, 0_L) - \frac{\sigma(\bullet, +1_L) + \sigma(\bullet, -1_L)}{2} \right] = \frac{2 V_{0,0}^{UQ_0} + (\frac{1}{3} - \cos^2 \theta) V_{2,0}^{UQ_0}}{2 W_{0,0} + (\frac{1}{3} - \cos^2 \theta) W_{2,0}}, \\
A_{UQ_1} &= \frac{\sigma(\bullet, I_1) - \sigma(\bullet, I_3)}{2 \langle \sigma \rangle} = \frac{\sin \theta \cos \theta \sin \phi U_{2,1}^{UQ_1}}{2 W_{0,0} + (\frac{1}{3} - \cos^2 \theta) W_{2,0}}, \\
A_{UQ_2} &= \frac{1}{2 \langle \sigma \rangle} \left[\frac{\sigma(\bullet, \phi_B = 0) + \sigma(\bullet, \phi_B = \pi)}{2} - \frac{\sigma(\bullet, \phi_B = \pi/2) + \sigma(\bullet, \phi_B = 3\pi/2)}{2} \right] \\
&= \frac{\sin^2 \theta \cos 2\phi U_{2,2}^{UQ_2}}{2 W_{0,0} + (\frac{1}{3} - \cos^2 \theta) W_{2,0}}.
\end{aligned} \tag{5}$$

Here, θ and ϕ are polar and azimuthal angles of the ℓ^+ momentum. The solid circle indicates that the proton is unpolarized. The three tensor polarizations are illustrated in Fig.1.




FIGURE 1. Q_0 , Q_1 , and Q_2 polarizations [4].

The Q_0 polarization asymmetry is in principle the same as the one in defining the tensor structure function b_1 . On the other hand, Q_2 is associated with the tensor asymmetry in the transverse plane. The Q_1 polarization is very peculiar. The deuteron is polarized with the angle 45° or 135° with respect to the longitudinal axis, and the cross section difference is taken. This kind of peculiar asymmetry does not exist, of course, for the proton. This “intermediate” (according to Ref. [4]) polarization asymmetry could lead to a new unique field of spin physics. A parton-model analysis suggests that the asymmetry Q_0 should be related to b_1 type tensor distributions as [4]

$$A_{UQ_0} = \frac{\sum_a e_a^2 [f_1^a(x_1) \bar{b}_1^a(x_2) + \bar{f}_1^a(x_1) b_1^a(x_2)]}{\sum_a e_a^2 [f_1^a(x_1) \bar{f}_1^a(x_2) + \bar{f}_1^a(x_1) f_1^a(x_2)]}, \quad (6)$$

where f_1^a and b_1^a are unpolarized and tensor-polarized distributions for the quark a , and \bar{f}_1^a and \bar{b}_1^a are those for the antiquark. The momentum fractions are given by x_1 and x_2 in the proton and the deuteron. For example, b_1^a is given by $b_1^a = [q_a(0) - (q_a(+1) + q_a(-1))/2]/2$. We mentioned the sum rule $\int dx b_1(x) = 0$. However, this relation crucially depends on the antiquark tensor polarization \bar{b}_1^a , which cannot be well probed in the electron scattering. One of the major advantages of using the pd reaction is to find \bar{b}_1^a rather easily, at least according to the formalism. The situation is similar to the case that the detailed x dependence of \bar{u}/\bar{d} was clarified by the Fermilab-E866 Drell-Yan experiment although the difference was originally indicated by the NMC data [3]. For example, in the large- x_F region, Eq.(6) becomes

$$A_{UQ_0}(\text{large } x_F) \approx \frac{\sum_a e_a^2 f_1^a(x_1) \bar{b}_1^a(x_2)}{\sum_a e_a^2 f_1^a(x_1) \bar{f}_1^a(x_2)}. \quad (7)$$

Therefore, the antiquark tensor distributions \bar{b}_1^a can be determined if the unpolarized distributions are well known in the proton and deuteron.

Another advantage of the pd Drell-Yan is that the \bar{u}/\bar{d} asymmetry can be investigated for the transversity distributions [5]. Because they have chiral-odd property,

they cannot be investigated in the inclusive electron scattering. The pd/pp Drell-Yan ratio is given by [5]

$$R_{pd} \equiv \frac{\Delta_{(T)}\sigma_{pd}}{2\Delta_{(T)}\sigma_{pp}} = \frac{\sum_a e_a^2 [\Delta_{(T)}q_a(x_1)\Delta_{(T)}\bar{q}_a^d(x_2) + \Delta_{(T)}\bar{q}_a(x_1)\Delta_{(T)}q_a^d(x_2)]}{2\sum_a e_a^2 [\Delta_{(T)}q_a(x_1)\Delta_{(T)}\bar{q}_a(x_2) + \Delta_{(T)}\bar{q}_a(x_1)\Delta_{(T)}q_a(x_2)]}$$

$$\rightarrow \frac{1}{2} \left[1 + \frac{\Delta_{(T)}\bar{d}(x_2)}{\Delta_{(T)}\bar{u}(x_2)} \right]_{x_2 \rightarrow 0} \quad \text{for } x_F \rightarrow 1, \quad (8)$$

where $\Delta_{(T)} = \Delta$ or Δ_T depending on the longitudinal or transverse polarization, and $\Delta_{(T)}\sigma$ indicates the polarized cross-section difference. In the second line, $x_F \rightarrow 1$ limit is taken together with the assumption $\Delta_{(T)}u_v(x \rightarrow 1) \gg \Delta_{(T)}d_v(x \rightarrow 1)$. Leading-order numerical results are shown in Fig.2, where $r_{\bar{q}} \equiv \Delta_{(T)}\bar{u}/\Delta_{(T)}\bar{d}$. It obviously indicates that the ratio is suitable for finding the polarized light antiquark distributions, particularly in the large- x_F region.

In this way, we found that the pd Drell-Yan is an alternative way of studying the deuteron spin structure to the electron scattering. Because other studies of spin-1 physics are not discussed in this paper, the author hopes that the interested reader will look at the papers in the extensive lists of Ref. [4,5].




FIGURE 2. pd/pp Drell-Yan ratio [5].

SUMMARY

We explained the possibilities of investigating the polarized deuteron. Exploring a new field of hadron spin, namely the tensor structure, we should be able to test our knowledge of high-energy spin physics. Because it has not been well investigated yet, we may encounter an unexpected result. Depending on the HERMES measurement in the near future, this kind of topic could become an interesting area of hadron physics. We also expressed the importance of polarized proton-deuteron reactions, which are possible not only in collider experiments but also in fixed-target ones.

REFERENCES

1. P. Hoodbhoy, R. L. Jaffe, and A. Manohar, *Nucl. Phys.* **B312**, 571 (1989).
2. F. E. Close and S. Kumano, *Phys. Rev.* **D42**, 2377 (1990).
3. S. Kumano, *Phys. Rep.* **303**, 183 (1998).
4. S. Hino and S. Kumano, *Phys. Rev.* **D59**, 094026 (1999); **D60**, 054018 (1999).
5. S. Kumano and M. Miyama, *Phys. Lett.* **B497**, 149 (2000).